

- effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med* 1996;335:1001-9.
14. Shepherd J, Cobbe SM, Ford I, Isles CG, Lorimer AR, MacFarlane PW, et al, for the West of Scotland Coronary Prevention Study Group. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med* 1995;333:1304-7.
 15. Scandinavian Simvastatin Survival Study Group. Randomized trial of cholesterol lowering in 4,444 patients with coronary artery disease (4S). *Lancet* 1994;344:1383-9.
 16. Downs JR, Clearfield M, Weis S, Whitney E, Shapiro DR, Beere PA, et al, for the AFCAPS/TextCAPS Research Group. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels. Results of AFCAPS/TextCAPS. *JAMA* 1998;279:1615-22.
 17. Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. The Long-term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group. *N Engl J Med* 1998;339:1349-57.
 18. Schwartz GG, Oliver M, Ezekowitz MD, Ganz P, Waters D, Kane JP, et al. Rationale and design of the Myocardial Ischemia Reduction with Aggressive Cholesterol Lowering (MIRACL) study that evaluates atorvastatin in unstable angina pectoris and in non-Q-wave acute myocardial infarction. *Am J Cardiol* 1998;81:578-81.
 19. Schwartz GG, Olsson AG, Ezekowitz MD, Ganz P, Oliver MF, Waters D, et al, Myocardial Ischemia Reduction with Aggressive Cholesterol Lowering (MIRACL) Study Investigators. Effects of atorvastatin on early recurrent ischemic events in coronary syndromes: a randomized controlled trial. *JAMA* 2001;285:1758-60.
 20. O'Driscoll G, Green D, Taylor RR. Simvastatin, an HMG-coenzyme A reductase inhibitor, improves endothelial function within 1 month. *Circulation* 1997;95:1126-31.
 21. Tsunekawa T, Hayashi T, Kano H, Sumi D, Matsui-Hirai H, Thakur NK, et al. Cerivastatin, a hydroxymethylglutaryl coenzyme A reductase inhibitor, improves endothelial function in elderly diabetic patients within 3 days. *Circulation* 2001;104:376-9.
 22. Tunstall-Pedoe H, Kuulasmaa K, Amouyel P, Arveiler D, Rajakangas AM, Pajak A. Myocardial infarction and coronary deaths in the World Health Organization MONICA Project. Registration procedures, event rates, and case-fatality rates in 38 populations from 21 countries on four continents. *Circulation* 1994;90:583-612.
 23. The Joint European Society of Cardiology/American College of Cardiology Committee. Myocardial infarction redefined: a consensus document of the Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of Myocardial Infarction. *J Am Coll Cardiol* 2000;36:959-69.
 24. Braunwald E, Antman EM, Beasley JW, Califf RM, Cheitlin MD, Hochman JS, et al. ACC/AHA 2002 guideline update for the management of patients with unstable angina and non-ST segment elevation myocardial infarction. *J Am Coll Cardiol* 2002;40:1366-74.
 25. Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomized placebo-controlled trial. *Lancet* 2002;360:7-22.

Submitted Sep 24, 2003; accepted Dec 11, 2003.

Available online Feb 23, 2004.

INVITED COMMENTARY

Janet T. Powell, MD, PhD

Few patients with peripheral arterial disease have a disease-free coronary circulation.¹ Reconstructive procedures impose further stress on the diseased coronary circulation, and cardiac morbidity and mortality are recognized complications of vascular surgery. The article by Durazzo et al in this issue of *The Journal* makes the exciting suggestion that a short course of perioperative statins could minimize these complications.

Before we all rush into new statin protocols, it is prudent to examine the evidence carefully. One hundred patients, scheduled for elective vascular procedures, were randomized to receive either 20 mg of atorvastatin or placebo, and the trial was designed to show a reduction in postoperative cardiovascular events from 22% to 1% within 6 months. This was a challenging target to set, because even in patients at the highest cardiac risk undergoing vascular surgery under general anaesthesia the perioperative cardiovascular event rate is 34%.³ In the trial by Durazzo et al,² patients in the placebo group underwent more procedures under spinal anaesthesia, 13 of 46 vs 5 of 44 in the atorvastatin group. Therefore the incidence of events in the placebo group (26%) was very high, unless spinal anaesthesia was used, because these patients were more likely to have American Society of Anesthesiologists grade III and IV disease. In contrast, statin treatment reduced the incidence of cardiovascular events to 8%, a result that just achieved significance $P = .031$. The excess events in the placebo group were predominantly myocardial infarction and stroke. Survival analysis showed a more convincing benefit in favor of atorvastatin treatment. Confirmation of these results from other trials is awaited eagerly.

So there is cautious optimism for an effective, simple, safe, new treatment to minimize perioperative cardiac morbidity and mortality after vascular surgery. Further support for this optimism comes from retrospective analyses of in-hospital mortality after vascular surgery and from late mortality after successful aortic

aneurysm repair.^{4,5} Statin therapy was more common in patients discharged after vascular surgery than in those who died in hospital (25% vs 8%; $P < .001$).⁴ Late cardiovascular deaths after aneurysm repair were reported in 9% of statin users compared with 38% in nonusers.⁵ All of these studies reported that the benefit of statins is independent of the use of β -blockers.

How do statins produce these effects? Reduction of serum lipid concentrations is an unlikely explanation for immediate beneficial effects of statins on perioperative morbidity and mortality. Two studies started with the hypothesis that statins exert their beneficial perioperative effect by plaque stabilization.^{2,4} Statin therapy also reduces postprocedural myocardial infarction and death after percutaneous coronary intervention.⁶ In this context, analysis of the serum C-reactive protein concentration indicated that statins exert their effect by an anti-inflammatory mechanism, statin therapy having most marked benefit in patients with the highest preprocedural C-reactive protein concentrations.⁶

It is disappointing that only a minority of patients (15%) in the trial by Durazzo et al² continued to receive statin therapy after 45 days. Vascular surgeons must not ignore the mounting evidence that statins benefit the cardiovascular health and survival of all patients with peripheral arterial disease.⁷

REFERENCES

1. Hertzner NR, Beven EG, Young JR, O'Hara PJ, Ruschhaupt WF III, Graor RA, et al. Coronary artery disease in peripheral vascular patients: a classification of 1000 coronary angiograms and results of surgical management. *Ann Surg* 1984;199:223-33.
2. Poldermans D, Boersma E, Bax JJ, Thomson IR, Van De Ven LLM, Blankensteijn JD, et al, for the Dutch Echocardiographic Cardiac Risk Evaluation Applying Stress Echocardiography Study Group. The effect of

- bisoprolol on perioperative mortality and myocardial infarction in high risk patients undergoing vascular surgery. *N Engl J Med* 1999;341:1789-94.
3. Poldermans D, Bax JJ, Kertai MD, Krenning B, Westerhout CM, Schinkel AFL, et al. Statins are associated with a reduced incidence of perioperative mortality in patients undergoing major noncardiac vascular surgery. *Circulation* 2003;107:1848-51.
 4. Kertai MD, Boersma E, Westerhout CM, van Domburg R, Klein J, Bax JJ, et al. Association between long-term statin use and mortality after successful abdominal aortic aneurysm surgery. *Am J Med* 2004;116:96-103.
 5. Chan AW, Bhatt DL, Chew DP, Reginelli J, Schneider JP, Topol EJ, et al. Relation of inflammation and benefit of statins after percutaneous coronary interventions. *Circulation* 2003;107:1750-6.
 6. Hiatt HR. Pharmacologic therapy for peripheral arterial disease and claudication. *J Vasc Surg* 2002;36:1283-91.



We have the answers
you are looking for.



Visit us at:

<http://www.vascularweb.org>